

Non-Abelian T-duality in Pre-Big-Bang Cosmology

A. Bossard¹ and N. Mohammadi²

*Laboratoire de Mathématiques et Physique Théorique³
Université François Rabelais
Faculté des Sciences et Techniques
Parc de Grandmont
F-37200 Tours - France.*

Abstract

We study the impact of non-Abelian T-duality transformations on a string based cosmological model. The implementation of the pre-big-bang scenario is investigated. We found a region of the dual phase where such a picture is possible.

¹e-mail: bossard@celfi.phys.univ-tours.fr

²e-mail: nouri@celfi.phys.univ-tours.fr

³CNRS UPRES-A 6083

1 Introduction

One of the most impressive theoretical constructions of the past century is undoubtedly the standard cosmological model based on the general relativistic Friedmann-Lemaître-Robertson-Walker (FLRW) metric. Not only it does explain the Hubble law, but it also led Gamov to predict the presence of a cosmic microwave background (CMB) radiation, detected many years later. Despite its many successes, such as the explanation of nucleosynthesis, this model is not exempt from drawbacks. Indeed, the standard scenario cannot satisfactorily explain the observed flatness of space-time nor the homogeneity of the CMB (the horizon problem) or the generation of cosmological perturbations responsible for the big structures of the universe. In order to remedy to these problems one is forced to assume an era of formidable expansion of the scale factor of the metric. This is the inflation scenario. However, as argued in [1], such an inflationary phase cannot eliminate the initial big-bang singularity of standard cosmology.

In order to deal with such a singularity one might need a quantum description of gravity. Presently, string theory seems to be the best candidate for describing the gravitational effects of matter at very small scales. In this theory, fundamental particles are described by tiny one dimensional strings embedded in external background fields. These backgrounds are not chosen arbitrarily. The internal coherence of string theory imposes constraints on the background fields, in particular the metric of space-time. Indeed, one demands that the two dimensional non linear sigma model description of the propagation of strings in these backgrounds should be invariant under Weyl transformations. This leads to the vanishing of the so called Weyl anomaly functions (or the generalized beta functions). This requirement results in a generalization of Einstein's equations of general relativity. Moreover, these Weyl anomaly equations can be viewed as equations of motion of an action (the string effective action) containing the Einstein-Hilbert term. This action describes the low energy excitations of string theory (the metric, the dilaton and the antisymmetric tensor).

Another important property of string theory is a set of transformations known as T-duality. These T-dualities relate seemingly two inequivalent sets of backgrounds. A string propagating in a space-time endowed with a first background would present the same dynamics when moving in another space-time endowed with a dual background. The natural area where the effects of these dualities might be observed is in cosmology. In seminal papers [2, 3], Veneziano and Gasperini have shown that a combination of Abelian T-duality, leading to an inversion of the scale factor of a flat FLRW metric (scale factor duality), together with a time reversal operation lead to a phase of inflation. This inflation is driven by a growing dilaton leading to a strong gravitational interaction. This phase (the dual phase) is valid for negative values of the time parameter and is called the pre-big-bang phase. However, the initial big-bang singularity is always present in this theory. It is sandwiched between two phases: the original flat FLRW solution and its Abelian dual.

Therefore, a suitable matching of the two phases must be found if one is to avoid the big-bang singularity. The answer to this problem could be found by including α' corrections in the string effective action (α' is the loop parameter of the two dimensional sigma model). The original analyses of Gasperini and Veneziano were carried out for the lowest order in α' . Higher order contributions in α' might be able to deal with the high curvature states typical of a big-bang singularity. The problem of regularization of

the big-bang singularity (the graceful exit problem) by means of higher α' contributions have been theoretically investigated in [4, 5, 6] and numerically in [7, 8, 9]. These studies indicate that the pre-big-bang cosmology could provide the necessary phase of inflation as well as a solution to the pre-big-bang singularity. This theory, however, still has its own drawbacks [10, 11].

In this paper, we would like to investigate the role of a more complicated T-duality, namely non-Abelian duality, in cosmology. Our aim is to figure out if a pre-big-bang picture can be implemented for a background corresponding to a non flat FLRW space-time. This space-time presents a natural setting for the application of non-Abelian T-duality. We are also motivated by the fact that the string effective action is invariant under non-Abelian T-duality transformations [12]. We are able to find a restricted region of the dual phase where a pre-big-bang approach does indeed apply. The difficulty in reaching a conclusion regarding the whole dual phase stems from the absence of any symmetries in the dual backgrounds. This situation is very typical of non-Abelian T-duality and is due to its non reversibility [13].

The paper is organized as follows: In section 2, we formulate the general problem and summarize the main results of non-Abelian T-duality. In particular, an expression relating the matter content in the original theory and its dual counterpart is given. This relation is essential for the extension of T-duality transformations in the presence of matter fields. We briefly recall, in section 3, the important features of the pre-big-bang cosmological model. Our example, based on a non flat FLRW metric, is studied in section 4. Finally, we present our conclusions and outlook in section 5.

2 String effective action and T-dualities

Our starting point is the following model as defined by the action

$$S = S_{\text{eff}} + S_{\text{mat}} , \quad (2.1)$$

where S_{mat} denotes the action for the matter content of some cosmological model and S_{eff} is the low energy effective action of string theory given by

$$S_{\text{eff}} = \int d^D X \sqrt{-G} e^{-\varphi} \left[R + (\partial\varphi)^2 - \frac{1}{12} H_{MNP} H^{MNP} + \Lambda \right] . \quad (2.2)$$

In this expression, G is the determinant of the metric G_{MN} and R is the associated Ricci scalar. The field φ is the usual dilaton and H_{MNP} , defined by $H_{MNP} = \partial_M B_{NP} + \partial_N B_{PM} + \partial_P B_{MN}$, is the torsion of the antisymmetric field B_{MN} . Finally, Λ stands for a possible cosmological constant. In the absence of matter fields, the equations of motion of the action S would reduce to the vanishing of the Weyl anomaly coefficients $\bar{\beta}^{(\omega)}$ defined by

$$\begin{aligned} \bar{\beta}_{MN}^{(G)} &\equiv R_{MN} + \nabla_M \partial_N \varphi - \frac{1}{4} H_{MPQ} H_N{}^{PQ} , \\ \bar{\beta}_{MN}^{(B)} &\equiv -\frac{1}{2} \nabla^P H_{MNP} + \frac{1}{2} H_{MNP} \partial^P \varphi , \\ \bar{\beta}^{(\varphi)} &\equiv -\frac{1}{4} \nabla^2 \varphi + \frac{1}{4} \partial_P \varphi \partial^P \varphi - \frac{1}{24} H_{MNP} H^{MNP} - \frac{\Lambda}{4} . \end{aligned} \quad (2.3)$$

In the presence of matter, however, the equations of motion of the action S can be cast in the following form

$$\begin{aligned}\bar{\beta}_{MN}^{(G)} &= \frac{1}{2}G_{MN}E^{(\varphi)} + e^\varphi T_{MN}^{(G)} , \\ \bar{\beta}_{MN}^{(B)} &= e^\varphi T_{MN}^{(B)} , \\ E^{(\varphi)} &= e^\varphi T^{(\varphi)} ,\end{aligned}\tag{2.4}$$

where

$$\begin{aligned}E^{(\varphi)} &\equiv \bar{\beta}_{PQ}^{(G)}G^{PQ} - 4\bar{\beta}^{(\varphi)} , \\ T_{MN}^{(G)} &\equiv -\frac{1}{\sqrt{-G}}\frac{\delta S_{\text{mat}}}{\delta G^{MN}} , \\ T_{MN}^{(B)} &\equiv G_{MP}G_{NQ}\frac{1}{\sqrt{-G}}\frac{\delta S_{\text{mat}}}{\delta B_{PQ}} , \\ T^{(\varphi)} &\equiv \frac{1}{\sqrt{-G}}\frac{\delta S_{\text{mat}}}{\delta \varphi} .\end{aligned}\tag{2.5}$$

We have allowed here the matter to depend on the dilaton as well as the antisymmetric tensor field. This is in order to have the most general setting for our scenario.

Let us now briefly recall the T-duality transformations present in the string effective action S_{eff} . The origin of these transformations resides in the two dimensional non-linear sigma model description of string theory [14, 15, 16]. We will consider here the simplest version of these transformations. In order to do so, the target space coordinates X^M are split as (x^μ, y^i) where we assume that an isometry group acts on the coordinates y^i only. The dependence on y^i of the metric G_{MN} and the antisymmetric tensor B_{MN} appears only through the one form $e_i^a dy^i$. In other words, our backgrounds are such that

$$\begin{aligned}G_{\mu\nu} &= G_{\mu\nu}(x) , \\ B_{\mu\nu} &= B_{\mu\nu}(x) , \\ G_{\mu i} &= Z_{\mu a}(x)e_i^a(y) , \\ B_{\mu i} &= W_{\mu a}(x)e_i^a(y) , \\ G_{ij} &= e_i^a(y)S_{ab}(x)e_j^b(y) , \\ B_{ij} &= e_i^a(y)v_{ab}(x)e_j^b(y) , \\ \varphi &= \varphi(x) ,\end{aligned}\tag{2.6}$$

Here, $e_i^a(y)$ are vielbeins satisfying

$$\partial_i e_j^a - \partial_j e_i^a = -f_{bc}^a e_i^b e_j^c\tag{2.7}$$

and f_{bc}^a are the structure constants of the non-Abelian isometry group acting on the string backgrounds.

With these assumptions, the string effective action S_{eff} remains invariant when the original backgrounds $\omega = (G_{MN}, B_{MN}, \varphi)$ are replaced by their duals $\tilde{\omega} = (\tilde{G}_{MN}, \tilde{B}_{MN}, \tilde{\varphi})$ [12]. The latter are given by

$$\tilde{G}_{\mu\nu} = G_{\mu\nu} - \bar{S}^{ab}(Z_{\mu a}Z_{\nu b} - W_{\mu a}W_{\nu b}) + \bar{A}^{ab}(Z_{\mu a}W_{\nu b} - W_{\mu a}Z_{\nu b})$$

$$\begin{aligned}
\tilde{B}_{\mu\nu} &= B_{\mu\nu} - \bar{A}^{ab}(Z_{\mu a}Z_{\nu b} - W_{\mu a}W_{\nu b}) + \bar{S}^{ab}(Z_{\mu a}W_{\nu b} - W_{\mu a}Z_{\nu b}) \\
\tilde{G}_{\mu i} &= -(Z_{\mu a}\bar{A}^{ab} + W_{\mu a}\bar{S}^{ab})\eta_{bi} \\
\tilde{B}_{\mu i} &= -(Z_{\mu a}\bar{S}^{ab} + W_{\mu a}\bar{A}^{ab})\eta_{bi} \\
\tilde{G}_{ij} &= \eta_{ia}\bar{S}^{ab}\eta_{bj} \\
\tilde{B}_{ij} &= \eta_{ia}\bar{A}^{ab}\eta_{bj} \\
\tilde{\varphi} &= \varphi - \ln(\det M_{ab}) ,
\end{aligned} \tag{2.8}$$

where η_{ia} is a unit matrix and E_a^i are the inverses of the vielbeins e_a^i . Moreover, $\bar{S}^{ab}$ and $\bar{A}^{ab}$ are the symmetric and antisymmetric parts of the inverse of

$$M_{ab} \equiv (S_{ab} + v_{ab}) + y^i \eta_{ic} f_{ab}^c . \tag{2.9}$$

The non-Abelian T-duality transformations lead also to some relations between the Weyl anomaly coefficients of the original backgrounds and the Weyl anomaly coefficients of the dual backgrounds⁴. These are given by [12]

$$\bar{\beta}^{(\tilde{\omega})} = \sum_{\omega} \frac{\delta \tilde{\omega}}{\delta \omega} \bar{\beta}^{(\omega)} . \tag{2.10}$$

It is remarkable that the duality transformations leave invariant the quantity $E^{(\varphi)}$ [12]. Namely,

$$E^{(\tilde{\varphi})} = E^{(\varphi)} . \tag{2.11}$$

In this paper we will consider exclusively bloc-diagonal backgrounds ($Z_{\mu a} = W_{\mu a} = 0$). The calculation of the partial derivatives $\delta \tilde{\omega} / \delta \omega$, given at the end of ref. [12], shows that, in a bloc-diagonal case, the previous relations (2.10) take the form

$$\begin{aligned}
\bar{\beta}_{MN}^{(\tilde{G})} &= \frac{\delta \tilde{G}_{MN}}{\delta G_{PQ}} \bar{\beta}_{PQ}^{(G)} + \frac{\delta \tilde{G}_{MN}}{\delta B_{PQ}} \bar{\beta}_{PQ}^{(B)} , \\
\bar{\beta}_{MN}^{(\tilde{B})} &= \frac{\delta \tilde{B}_{MN}}{\delta G_{PQ}} \bar{\beta}_{PQ}^{(G)} + \frac{\delta \tilde{B}_{MN}}{\delta B_{PQ}} \bar{\beta}_{PQ}^{(B)} .
\end{aligned} \tag{2.12}$$

Our aim here is to study, for a simple model, non-Abelian duality transformations in the presence of matter fields. The demand that these transformations are still a symmetry of the full action S will require the original matter tensors $T^{(\omega)}$ to have dual counterparts $T^{(\tilde{\omega})}$. Indeed, using the equations of motion of the original and dual models together with the relations in (2.11, 2.12), one can show that

$$\begin{aligned}
e^{\tilde{\varphi}} T^{(\tilde{\varphi})} &= e^{\varphi} T^{(\varphi)} , \\
e^{\tilde{\varphi}} T_{MN}^{(\tilde{G})} &= e^{\varphi} \left[\frac{1}{2} \left(\frac{\delta \tilde{G}_{MN}}{\delta G_{PQ}} G_{PQ} - \tilde{G}_{MN} \right) T^{(\varphi)} + \frac{\delta \tilde{G}_{MN}}{\delta G_{PQ}} T_{PQ}^{(G)} + \frac{\delta \tilde{G}_{MN}}{\delta B_{PQ}} T_{PQ}^{(B)} \right] , \\
e^{\tilde{\varphi}} T_{MN}^{(\tilde{B})} &= e^{\varphi} \left[\frac{1}{2} \frac{\delta \tilde{B}_{MN}}{\delta G_{PQ}} G_{PQ} T^{(\varphi)} + \frac{\delta \tilde{B}_{MN}}{\delta G_{PQ}} T_{PQ}^{(G)} + \frac{\delta \tilde{B}_{MN}}{\delta B_{PQ}} T_{PQ}^{(B)} \right] .
\end{aligned} \tag{2.13}$$

The matter sector in the dual theory will be determined for our cosmological model. Before tackling this issue, we will start by a short review of pre-big-bang cosmology.

⁴The equations of motion and the Weyl anomaly functions for the dual theory are obtained from those of the original model by replacing the untilded quantities by tilded ones.

3 Abelian pre-big-bang cosmology

The pre-big-bang picture was put forward by Veneziano and Gasperini [3, 17, 18, 19]. Its main ingredient is the use of Abelian T-duality in a string cosmological model. The space-time is four dimensional with coordinates $X^M = (t, y^i)$ where $y^i = (x, y, z)$. The original backgrounds are chosen to be given by

$$\begin{aligned} G_{MN} &= \text{diag}(-1, a^2(t), a^2(t), a^2(t)) , \\ B_{MN} &= 0 , \\ \varphi &= \varphi(t) . \end{aligned} \tag{3.1}$$

The non-vanishing components of the Weyl anomaly coefficients $\bar{\beta}^{(\omega)}$ are

$$\begin{aligned} \bar{\beta}_{00}^{(G)} &= \ddot{\varphi} - 3\dot{H} - 3H^2 , \\ \bar{\beta}_{ij}^{(G)} &= \delta_{ij}a^2[\dot{H} + 3H^2 - H\dot{\varphi}] , \end{aligned} \tag{3.2}$$

and

$$E^{(\varphi)} = 12H^2 + 6\dot{H} - 2\ddot{\varphi} - 6H\dot{\varphi} + \dot{\varphi}^2 + \Lambda . \tag{3.3}$$

The Hubble function is defined by the usual relation $H(t) \equiv \frac{d}{dt} \ln a(t)$.

The coupling of the matter in this model is that of a perfect fluid, as in standard cosmology. The matter tensors $T_{MN}^{(\omega)}$ are such that

$$\begin{aligned} T_{MN}^{(G)} &= \text{diag}(\rho, pa^2, pa^2, pa^2) , \\ T_{MN}^{(B)} &= 0 , \\ T^{(\varphi)} &= 0 . \end{aligned} \tag{3.4}$$

The equations of motion corresponding to the action S are

$$\begin{aligned} \ddot{\varphi} - 3\dot{H} - 3H^2 &= e^\varphi \rho , \\ \dot{H} + 3H^2 - H\dot{\varphi} &= e^\varphi p , \\ 12H^2 + 6\dot{H} - 2\ddot{\varphi} - 6H\dot{\varphi} + \dot{\varphi}^2 + \Lambda &= 0 . \end{aligned} \tag{3.5}$$

The original backgrounds in (3.1) possess three translation symmetries (Abelian isometries). The application of Abelian T-duality transformations [14] leads to the following dual backgrounds

$$\begin{aligned} \tilde{G}_{MN} &= \text{diag}(-1, a^{-2}(t), a^{-2}(t), a^{-2}(t)) , \\ \tilde{B}_{MN} &= 0 , \\ \tilde{\varphi} &= \varphi(t) - 6 \ln a(t) \end{aligned} \tag{3.6}$$

The corresponding non-vanishing dual Weyl anomaly coefficients $\bar{\beta}^{(\tilde{\omega})}$ are as follows

$$\begin{aligned} \bar{\beta}_{00}^{(\tilde{G})} &= \ddot{\varphi} - 3\dot{H} - 3H^2 , \\ \bar{\beta}_{ij}^{(\tilde{G})} &= -\delta_{ij}a^{-2}[\dot{H} + 3H^2 - H\dot{\varphi}] , \end{aligned} \tag{3.7}$$

and

$$E^{(\tilde{\varphi})} = 12H^2 + 6\dot{H} - 2\ddot{\varphi} - 6H\dot{\varphi} + \dot{\varphi}^2 + \Lambda . \quad (3.8)$$

If we now suppose, in analogy with the original model, that the matter fields in the dual model are such that

$$\begin{aligned} T^{(\tilde{\varphi})} &= 0 , \\ T_{MN}^{(\tilde{B})} &= 0 , \\ T_{MN}^{(\tilde{G})} &= \text{diag}(\tilde{\rho}, \tilde{p}a^{-2}, \tilde{p}a^{-2}, \tilde{p}a^{-2}) , \end{aligned} \quad (3.9)$$

then the equations of motion of the dual theory are

$$\begin{aligned} \ddot{\varphi} - 3\dot{H} - 3H^2 &= e^{\tilde{\varphi}} \tilde{\rho} , \\ \dot{H} + 3H^2 - H\dot{\varphi} &= -e^{\tilde{\varphi}} \tilde{p} , \\ 12H^2 + 6\dot{H} - 2\ddot{\varphi} - 6H\dot{\varphi} + \dot{\varphi}^2 + \Lambda &= 0 . \end{aligned} \quad (3.10)$$

These equations are the same as the ones in (3.5) provided that the matter fields in the original model and its dual are related by

$$\begin{aligned} e^{\varphi} \rho &= e^{\tilde{\varphi}} \tilde{\rho} , \\ e^{\varphi} p &= -e^{\tilde{\varphi}} \tilde{p} . \end{aligned} \quad (3.11)$$

Therefore, the duality transformations of S_{eff} can be extended to the full action $S = S_{\text{eff}} + S_{\text{mat}}$. Notice also that if the original matter obeys the equation of state $\rho = \alpha p$, where α is a constant, then the matter in the dual theory satisfies the equation of state $\tilde{\rho} = -\alpha \tilde{p}$. We deduce that if α is positive then the pressure in the dual phase must be negative.

Let us now examine the cosmological effects of Abelian T-duality. We suppose, for simplicity, that the dilaton field is constant $\varphi = \varphi_0$ and the cosmological constant vanishes. In this case, the last equation of (3.5) leads to $a(t) \sim \sqrt{t}$ while the two others yield the equation of state $\rho = 3p$. Hence the universe is in a radiation dominated era. The energy density ρ is related to the scale factor a by $\rho \sim a^{-4}$. An important property of this standard cosmological solution is that of not being defined for all values of t and has a big-bang singularity at $t = 0$.

The idea behind pre-big-bang cosmology is to seek a solution which is defined for all values of t and avoids the big-bang singularity. This solution is found through the use of T-duality transformations combined with a time reversal operation. Indeed, the equations of motion of the original model (3.5) as well as those of the dual theory (3.10) are form invariant under $t \rightarrow -t$.

The pre-big-bang picture can be summarized by the graph in Figure 1.

One starts from the original solution (region I) as defined by

$$\begin{aligned} a &\sim \sqrt{t} , \\ \rho &= 3p \sim a^2 , \\ \varphi &= \varphi_0 \end{aligned} \quad (3.12)$$




Figure 1: Abelian T-duality and time reversal

and constructs region II by applying a T-duality transformation. Region VI is reached by a time reversal operation on the solution of region II. Therefore, one obtains for region IV the following solution

$$\begin{aligned}
\tilde{a} &\sim \frac{1}{\sqrt{-t}} , \\
\tilde{\rho} &= -3\tilde{p} \sim \tilde{a}^{-2} , \\
\tilde{\varphi} &= \varphi_0 - 3 \ln(-t); .
\end{aligned} \tag{3.13}$$

This solution is defined for negative times and a full account of its characteristics can be found in [3, 17, 19, 18].

Gasperini and Veneziano suggested that by taking region IV when t is negative and region I when t is positive, one might avoid the big-bang singularity. However, this requires a non trivial mechanism for matching these two branches. We refer the reader to [4, 5, 6, 7, 8, 9] for more details on this subject.

4 Non-Abelian T-duality in pre-big-bang cosmology

Our goal in this article is to apply non-Abelian T-duality transformations to a cosmological model with a pre-big-bang scenario in sight. This is motivated by the fact that most of the interesting cosmological metrics present, in a natural way, non-Abelian isometries. We therefore consider a four dimensional FLRW space-time described by the coordinates $X^M = (t, y^i)$ with spherical spatial coordinates $y^i = (R, \Theta, \Phi)$. The string metric is described by

$$G_{MN} = \text{diag} \left(-1, \frac{a^2(t)}{1 - kR^2}, a^2(t)R^2, a^2(t)R^2 \sin^2(\Theta) \right) . \tag{4.1}$$

The other string background fields are given by

$$B_{MN} = 0 , \quad \varphi = \varphi_0 . \tag{4.2}$$

Notice that the spatially flat case, corresponding to $k = 0$, is the one treated in the previous section. We will assume from now on that k is different from zero.

In order to find the dual backgrounds, one has to cast the metric in a form suitable for the application of the formalism of non-Abelian T-duality. Therefore, one must determine the vielbeins of the spatial metric. We distinguish two cases. The first corresponds to k negative. This case is identified with the isotropic Bianchi V space. The Lie algebra of the isometry transformations of Bianchi V are known to have tracefull structure constants $f_{ab}^a \neq 0$. It is precisely for this kind of Lie algebras that non-Abelian T-duality does not apply. This problem with Bianchi V was first revealed in [20] and further examined in [21, 22, 23, 12]. For this reason, this case will not be considered any further.

The second case deals with k positive. This corresponds to the isotropic Bianchi IX space whose isometry Lie algebra is $SU(2)$ with $f_{ab}^a = 0$. Since there is only one maximally symmetric three dimensional metric with a constant positive curvature, a change of coordinates can be found [24] to bring the metric in (4.1) into the form

$$\begin{aligned} ds^2 &= -dt^2 + h(t) \text{Tr} \left[\left(g^{-1} dg \right) \left(g^{-1} dg \right) \right] , \\ &= -dt^2 + h(t) \delta_{ab} e_i^a e_j^b du^i du^j , \\ &= -dt^2 + h(t) \left(du^2 + dv^2 + dw^2 + 2 \cos(v) du dw \right) . \end{aligned} \quad (4.3)$$

In this expression, g is an element of the Lie group $SU(2)$. It is parametrized by $g = e^{uT_3} e^{vT_1} e^{wT_3}$, where T_a are related to the usual Pauli matrices σ_a by $T_a = \sigma_a/2i$. The function $h(t)$ is defined by $h(t) = a^2(t)/4k$ and the vielbeins are given by

$$e_i^a(y) = \begin{pmatrix} \sin(v) \sin(w) & \cos(w) & 0 \\ \sin(v) \cos(w) & -\sin(w) & 0 \\ \cos(v) & 0 & 1 \end{pmatrix} . \quad (4.4)$$

The indices a and i denote, respectively, the lines and rows of e_i^a .

To simplify matters, we assume a vanishing cosmological constant ($\Lambda = 0$). These backgrounds leads to

$$E^{(\varphi)} = 6 \left[\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + \frac{k}{a^2} \right] . \quad (4.5)$$

The other non-vanishing Weyl anomaly function $\bar{\beta}^{(\omega)}$ are

$$\begin{aligned} \bar{\beta}_{00}^{(G)} &= -3 \frac{\ddot{a}}{a} , \\ \bar{\beta}_{ij}^{(G)} &= G_{ij} F(t) , \\ F(t) &\equiv \ddot{a}/a + 2 (\dot{a}/a)^2 + 2k/a^2 . \end{aligned} \quad (4.6)$$

We also take the matter present in the universe to be described by the energy-momentum tensor

$$T_{MN}^{(G)} = \begin{pmatrix} \rho & 0 \\ 0 & p G_{ij} \end{pmatrix} , \quad T^{(\varphi)} = T_{MN}^{(B)} = 0 . \quad (4.7)$$




Figure 2: Big-bang and big-crunch singularities

The equations of motion are then

$$\begin{aligned} E^{(\varphi)} &= 0, \\ -3\frac{\ddot{a}}{a} &= e^{\varphi_0}\rho, \\ F(t) &= e^{\varphi_0}p, \end{aligned} \tag{4.8}$$

Combining these three equations leads to the equation of states $\rho = 3p$. Hence, as in the Abelian case, the matter is in a radiation form. One can then show that $C = (e^{\varphi_0}\rho a^4)/3$ is constant in time and the general solution for the equations of motion is given by

$$a(t) = \sqrt{C'} \left[1 - \left(1 - \frac{t}{\sqrt{C'}} \right)^2 \right]^{\frac{1}{2}}, \tag{4.9}$$

where $C' = C/k^2$. The graph of $a(t)$ is shown in Figure 2. This solution has a big-bang singularity at $t = 0$ and a big-crunch singularity at $t = 2\sqrt{C'}$.

Now we are in a position to construct the dual backgrounds as explained in the second section of this paper. One obtains, after a coordinate transformation from (t, u, v, w) to (t, r, θ, ϕ) [25], the following dual backgrounds⁵

$$\begin{aligned} \tilde{G}_{MN} &= \text{diag} \left(-1, \frac{1}{h(t)}, \frac{h(t)r^2}{\Delta}, \frac{h(t)r^2 \sin^2(\theta)}{\Delta} \right), \\ \tilde{B}_{MN} &= \frac{r^3 \sin(\theta)}{\Delta} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \\ \tilde{\varphi} &= \varphi(t) - \ln(h\Delta), \end{aligned} \tag{4.10}$$

⁵The change of variables is given by $u = r \sin(\theta) \cos(\phi)$, $v = r \sin(\theta) \sin(\phi)$, $w = r \cos(\theta)$.

where $\Delta = h^2(t) + r^2$. One can see that the metric of this dual backgrounds is still isotropic, due to the presence of the element $d\theta^2 + \sin^2(\theta)d\phi^2$ in the expression of ds^2 . But it is not homogeneous anymore. This kind of metric must then have a *center*.

The non-vanishing Weyl anomaly coefficients $\bar{\beta}^{(\omega)}$ of these dual backgrounds are

$$\begin{aligned}
E^{(\bar{\varphi})} &= E^{(\varphi)} , \\
\bar{\beta}_{23}^{(\bar{B})} &= 2r^3 \sin(\theta) \frac{h^2}{\Delta^2} F(t) , \\
\bar{\beta}_{00}^{(\tilde{G})} &= \frac{3C}{(4k)^2} \frac{1}{h^2} , \\
\bar{\beta}_{11}^{(\tilde{G})} &= -\frac{F(t)}{h} , \\
\bar{\beta}_{22}^{(\tilde{G})} &= -\frac{r^2 h (h^2 - r^2)}{\Delta^2} F(t) , \\
\bar{\beta}_{33}^{(\tilde{G})} &= -\frac{r^2 \sin^2(\theta) h (h^2 - r^2)}{\Delta^2} F(t) ,
\end{aligned} \tag{4.11}$$

By injecting these expressions in the equations of motion of the dual theory, we are led to the following form for the dual energy-momentum tensor

$$T_{MN}^{(\tilde{G})} = \begin{pmatrix} \tilde{\rho} & 0 & 0 & 0 \\ 0 & \tilde{p}_1 \tilde{G}_{11} & 0 & 0 \\ 0 & 0 & \tilde{p}_2 \tilde{G}_{22} & 0 \\ 0 & 0 & 0 & \tilde{p}_3 \tilde{G}_{33} \end{pmatrix} . \tag{4.12}$$

An explicit calculation yields the following expressions for the energy density and pressures appearing in $T_{MN}^{(\tilde{G})}$

$$\begin{aligned}
\tilde{\rho} &= 3K \frac{h^2 + r^2}{h} , \\
\tilde{p}_1 &= -K \frac{h^2 + r^2}{h} , \\
\tilde{p}_2 &= -K \frac{h^2 - r^2}{h} , \\
\tilde{p}_3 &= \tilde{p}_2 ,
\end{aligned} \tag{4.13}$$

where K is defined by $K = e^{-\varphi_0} C / (4k)^2$. Furthermore, the equations of motion of the dual theory are fulfilled only if

$$\begin{aligned}
T^{(\bar{\varphi})} &= 0 , \\
T_{23}^{(\bar{B})} &= -2K \tilde{B}_{23} h .
\end{aligned} \tag{4.14}$$

This means that in the dual phase, the matter depends on the dual antisymmetric tensor $\tilde{B}_{MN}$. This is to be contrasted with the original theory where we have set $B_{MN} = 0$ (implying $T_{MN}^{(B)} = 0$). The non-vanishing of $\tilde{B}_{MN}$ and $T_{MN}^{(\tilde{B})}$ breaks the isotropy observed in the dual metric $\tilde{G}_{MN}$. We should mention that the expressions of $T_{MN}^{(\tilde{G})}$, $T_{MN}^{(\tilde{B})}$ and $T^{(\bar{\varphi})}$ are compatible with the formal relations in (2.13).

5 Discussion

Let us now investigate whether one can set a pre-big-bang scenario based on the original backgrounds and their duals. As before, the equations of motion are still form invariant under time reversal $t \rightarrow -t$. Our original solution has two singularities and the pre-big-bang picture, based on non-Abelian duality and time reversal transformations, would be expected to avoid these singularities. This task, however, is difficult to achieve due to the fact that the dual backgrounds do not carry the symmetries of their original counterparts. Nonetheless, some conclusions can be reached in some limiting procedure. Two cases, depending on the relative values of r^2 and h^2 , emerge naturally in the study of the dual backgrounds.

The first of these two cases corresponds to the situation when $r^2 \ll h^2$. The dual backgrounds tend then to the following expressions

$$\begin{aligned}\tilde{G}_{MN} &\rightarrow \text{diag}(-1, \frac{1}{h}, \frac{r^2}{h}, \frac{r^2 \sin^2(\theta)}{h}), \\ \tilde{B}_{MN} &\rightarrow 0, \\ \tilde{\varphi} &\rightarrow \varphi_0 - 3 \ln h.\end{aligned}\tag{5.1}$$

Similarly, the dual matter fields are described by

$$\begin{aligned}\tilde{\rho} &\rightarrow 3Kh, \\ \tilde{p}_1 &\rightarrow \tilde{p} = -Kh, \\ \tilde{p}_2 &\rightarrow \tilde{p} = -Kh, \\ \tilde{p}_3 &\rightarrow \tilde{p} = -Kh, \\ T_{23}^{(\tilde{B})} &\rightarrow 0.\end{aligned}\tag{5.2}$$

In this limiting zone, the universe is described by a flat FLRW metric with a scale factor given by $1/\sqrt{h(t)}$. Furthermore, the only matter present is that of a perfect fluid obeying the equation of state $\tilde{\rho} = -3\tilde{p}$. One can, therefore, define a Hubble function $\tilde{H} = -\frac{d}{dt} \ln \sqrt{h}$.

In analogy with the Abelian pre-big-bang scenario, we construct in this limit a similar setting as represented in Figure 3. The Hubble function of region I is that of the original theory, while region IV corresponds to the metric in (5.1) but for negative t .

As one can see, the behaviors of $H(t)$ and $\tilde{H}(-t)$ are similar to what happened in the Abelian case (see Fig.1) near the big-bang singularity. Here again, one must solve the graceful exit problem (the matching of the two branches). There are, however, two other singularities (big-crunch) when $t \rightarrow \pm 2\sqrt{C'}$ (see Fig. 2). We speculate that higher order corrections in α' might cure the model of all these singularities.

The other limiting case corresponds to $r^2 \gg h^2$. Although, we are not able to construct a pre-big-bang cosmology for this limit, its study reveals some interesting features. The dual backgrounds in this zone tend to

$$\begin{aligned}\tilde{G}_{MN} &\rightarrow \text{diag}(-1, \frac{1}{h}, h, h \sin^2(\theta)), \\ \tilde{B}_{MN} &\rightarrow r \sin(\theta), \\ \tilde{\varphi} &\rightarrow \varphi_0 - \ln(hr^2).\end{aligned}\tag{5.3}$$




Figure 3: Non-Abelian T-duality and time reversal

On the other hand, the dual matter is represented by the quantities

$$\begin{aligned}
 \tilde{\rho} &\rightarrow 3K \frac{r^2}{h} , \\
 \tilde{p}_1 &\rightarrow -K \frac{r^2}{h} , \\
 \tilde{p}_2 &\rightarrow K \frac{r^2}{h} , \\
 \tilde{p}_3 &\rightarrow K \frac{r^2}{h} , \\
 T_{23}^{(\tilde{B})} &\rightarrow -2Khr \sin(\theta) .
 \end{aligned} \tag{5.4}$$

We notice that the r coordinate gets a special status in these backgrounds. The dual metric contains a sphere of radius $\sqrt{h}$ and whose surface is described by the coordinates θ and ϕ . When h is big the r direction shrinks while the sphere expands. The opposite happens for small h . This situation is similar to the procedure of dynamical compactification encountered in Kaluza-Klein cosmologies [26]. Furthermore, the pressure along the r direction is negative.

In conclusion, we have managed to put forward a pre-big-bang cosmological scenario in a very particular region ($r^2 \ll h^2$) in the dual phase. However, a global study of the full dual phase seems at the moment out of reach. The lack of symmetries in the dual backgrounds (absence of isotropy or homogeneity) does not allow a construction of a Hubble function. Therefore, a comparison with the original FLRW backgrounds is not, in general, possible.

Finally, we should mention that we have analysed another cosmological model based on Poisson-Lie T-duality. Unfortunately, the dual backgrounds are even more involved and do not reveal any known features. This shows all the difficulties whenever one deviates from the flat case of Abelian T-duality (scale factor duality). As already remarked in [27, 28], this is very puzzling from a string point of view: a fundamental string does not

see any differences between two backgrounds related by any T-duality transformations. Therefore, the role of isotropy and homogeneity in string cosmology is certainly worth investigating.

References

- [1] A. Borde and A. Vilenkin, “Eternal inflation and the initial singularity” *Phys. Rev. Lett.* **72**, 3305 (1994), gr-qc/9312022.
- [2] G. Veneziano, “Scale factor duality for classical and quantum strings,” *Phys. Lett. B* **265** (1991) 287.
- [3] M. Gasperini and G. Veneziano, “Pre - big bang in string cosmology,” *Astropart. Phys.* **1** (1993) 317 [arXiv:hep-th/9211021].
- [4] M. Gasperini, J. Maharana and G. Veneziano, “Graceful exit in quantum string cosmology,” *Nucl. Phys. B* **472** (1996) 349, hep-th/9602087.
- [5] M. Gasperini, M. Maggiore and G. Veneziano, “Towards a non-singular pre-big bang cosmology,” *Nucl. Phys. B* **494** (1997) 315, hep-th/9611039.
- [6] D. A. Easson and R. H. Brandenberger, “Nonsingular dilaton cosmology in the string frame,” *JHEP* **9909** (1999) 003, hep-th/9905175.
- [7] J. Maharana, E. Onofri and G. Veneziano, “A numerical simulation of pre-big bang cosmology,” *JHEP* **9804** (1998) 004, gr-qc/9802001.
- [8] T. Chiba, “Numerical study of inhomogeneous pre-big-bang inflationary cosmology,” *Phys. Rev. D* **59** (1999) 083508, gr-qc/9902003.
- [9] G. F. Ellis, D. C. Roberts, D. Solomons and P. K. Dunsby, “Solution To The Graceful Exit Problem In Pre-Big-Bang Cosmology,” *Phys. Rev. D* **62** (2000) 084004.
- [10] R. Brustein and G. Veneziano, “The Graceful exit problem in string cosmology,” *Phys. Lett. B* **329** (1994) 429 [arXiv:hep-th/9403060].
- [11] M. S. Turner and E. J. Weinberg, “Pre-big-bang inflation requires fine tuning,” *Phys. Rev. D* **56** (1997) 4604 [arXiv:hep-th/9705035].
- [12] A. Bossard and N. Mohammadi, “Non-Abelian duality in the string effective action,” *Nucl. Phys. B* **595** (2001) 93 [arXiv:hep-th/0008062].
- [13] A. Giveon and M. Roček, “On nonAbelian duality,” *Nucl. Phys. B* **421** (1994) 173 [arXiv:hep-th/9308154].
- [14] T. H. Buscher, “A Symmetry Of The String Background Field Equations,” *Phys. Lett. B* **194** (1987) 59.
- [15] M. Roček and E. Verlinde, “Duality, quotients, and currents,” *Nucl. Phys. B* **373** (1992) 630 [arXiv:hep-th/9110053].
- [16] X. C. de la Ossa and F. Quevedo, “Duality symmetries from nonAbelian isometries in string theory,” *Nucl. Phys. B* **403** (1993) 377 [arXiv:hep-th/9210021].
- [17] M. Gasperini, “Elementary introduction to pre-big bang cosmology and to the relic graviton background,” hep-th/9907067.

- [18] G. Veneziano, “A simple/short introduction to pre-big-bang physics/cosmology,” hep-th/9802057.
- [19] G. Veneziano, “String cosmology: The pre-big bang scenario,” hep-th/0002094.
- [20] M. Gasperini, R. Ricci and G. Veneziano, “A Problem with nonAbelian duality?,” Phys. Lett. B **319** (1993) 438 [arXiv:hep-th/9308112].
- [21] E. Tyurin, “On conformal properties of the dualized sigma models,” Phys. Lett. B **348** (1995) 386 [arXiv:hep-th/9411242].
- [22] E. Alvarez, L. Alvarez-Gaume and Y. Lozano, “On nonAbelian duality,” Nucl. Phys. B **424** (1994) 155 [arXiv:hep-th/9403155].
- [23] S. Elitzur, A. Giveon, E. Rabinovici, A. Schwimmer and G. Veneziano, “Remarks on nonAbelian duality,” Nucl. Phys. B **435** (1995) 147 [arXiv:hep-th/9409011].
- [24] S. Weinberg, “Gravitation and Cosmology,” Wiley, (1972).
- [25] J. Balog, P. Forgacs, N. Mohammedi, L. Palla and J. Schnittger, “On quantum T-duality in sigma models,” Nucl. Phys. B **535** (1998) 461 [arXiv:hep-th/9806068].
- [26] A. Chodos and S. Detweiler, “Where Has The Fifth-Dimension Gone?,” Phys. Rev. D **21** (1980) 2167.
- [27] A. Feinstein, R. Lazkoz and M. A. Vazquez-Mozo, “Closed inhomogeneous string cosmologies,” Phys. Rev. D **56** (1997) 5166, hep-th/9704173.
- [28] A. Feinstein, “Some aspects of pre big bang cosmology,” gr-qc/0106001.